



Risk factors associated with complications in lower-extremity reconstruction with the distally based sural flap: A systematic review and pooled analysis



Catherine de Blacam^a, Salih Colakoglu^b, Adeyemi A. Ogunleye^c,
John T. Nguyen^d, Ahmed M.S. Ibrahim^d, Samuel J. Lin^d,
Peter S. Kim^d, Bernard T. Lee^{d,*}

^a Plastic Surgery Higher Surgical Training Scheme, Royal College of Surgeons in Ireland, Dublin, Ireland

^b Department of Surgery, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

^c Department of Surgery, Harlem Hospital Center, Columbia University College of Physicians and Surgeons, New York, NY, USA

^d Department of Surgery, Division of Plastic and Reconstructive Surgery, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA

Received 14 May 2013; accepted 28 January 2014

KEYWORDS

Lower-extremity
reconstruction;
Sural flap;
Distally based sural
flap

Summary The distally based sural fasciocutaneous flap is one of the few options available for local flap reconstruction of soft-tissue defects in the lower one-third of the leg. Few studies have assessed risk factors associated with poor outcomes in this flap. A literature search was performed of MEDLINE, EMBASE, CINAHL and the Cochrane Library for articles evaluating the use of sural artery fasciocutaneous flaps for soft-tissue reconstruction of the leg. Data were pooled and a univariate analysis was performed to identify characteristics associated with increased morbidity. A logistic regression model was created, and odds ratios and p values associated with the development of complications were calculated. Sixty-one papers were identified which included data on 907 patients. The majority of sural flaps were used to cover defects of the heel (28.2%), foot (14.4%) or ankle (25.8%). Trauma was the most common indication, followed by ulcers and open fractures. Flap complications were recorded in 26.4% of cases with a flap loss rate of 3.2%. With multivariate analysis, venous insufficiency and increasing age were independent risk factors for complications. Patients with venous

* Corresponding author. Department of Surgery, Division of Plastic and Reconstructive Surgery, Beth Israel Deaconess Medical Center, Harvard Medical School, 110 Francis St., Suite 5A, Boston, MA 02215, USA. Tel.: +1 617 632 7835; fax: +1 617 632 7840.

E-mail address: blee3@bidmc.harvard.edu (B.T. Lee).