

The Utility of Outcome Studies in Plastic Surgery

Hani Sinno, MD, MEng*†
 Tassos Dionisopoulos, MD†
 Sumner A. Slavin, MD*
 Ahmed M. S. Ibrahim, MD*
 Kevin C. Chung, MD‡
 Samuel J. Lin, MD*

Summary: Outcome studies help provide the evidence-based science rationalizing treatment end results that factor the experience of patients and the impact on society. They improve the recognition of the shortcoming in clinical practice and provide the foundation for the development of gold standard care. With such evidence, health care practitioners can develop evidence-based justification for treatments and offer patients with superior informed consent for their treatment options. Furthermore, health care and insurance agencies can recognize improved cost-benefit options in the purpose of disease prevention and alleviation of its impact on the patient and society. Health care outcomes are ultimately measured by the treatment of disease, the reduction of symptoms, the normalization of laboratory results and physical measures, saving a life, and patient satisfaction. In this review, we outline the tools available to measure outcomes in plastic surgery and subsequently allow the objective measurements of plastic surgical conditions. Six major outcome categories are discussed: (1) functional measures; (2) preference-based measures and utility outcome scores; (3) patient satisfaction; (4) health outcomes and time; (5) other tools: patient-reported outcome measurement information system, BREAST-Q, and Tracking Operations and Outcomes for Plastic Surgeons; and (6) cost-effectiveness analysis. We use breast hypertrophy requiring breast reduction as an example throughout this review as a representative plastic surgical condition with multiple treatments available. (*Plast Reconstr Surg Glob Open* 2014;2:e189; doi: 10.1097/GOX.000000000000104; Published online 29 July 2014.)

Plastic surgeons have a substantial variation in surgical practice for similar diagnoses and conditions. For example, women with breast hypertrophy are offered breast reduction surgery with

over half a dozen different pedicles and incision types depending on the consultant surgeon and patient preference. This difference in practice is not limited to the health state/diagnosis, borders, race, ethics, or health care policy. Indeed, even within the same institution, patients with the same condition are treated differently with respect to medical and surgical practice. Whose technique is “superior”? Which patient is receiving “better care”? Does the lack of uniformity in surgical practice indicate that all techniques are equal for the care of one condition in different patients? Or does it mean that there is not enough evidence to prove which procedure is “superior”? Outcomes research helps answer such questions by providing the relation between clinical care and their end results.¹⁻³ As Clancy and Eisenberg⁴ outlined, outcomes research help measure the end results of health care.

*From the *Division of Plastic Surgery, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Mass.;*

†Division of Plastic Surgery, Sir Mortimer B. Davis Jewish General Hospital, McGill University, Montreal, Quebec, Canada; and ‡Division of Plastic Surgery, University of Michigan Medical School, Ann Arbor, Mich.

Received for publication December 27, 2012; accepted April 3, 2014.

Copyright © 2014 The Authors. Published by Lippincott Williams & Wilkins on behalf of The American Society of Plastic Surgeons. PRS Global Open is a publication of the American Society of Plastic Surgeons. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 3.0 License, where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially.

DOI: 10.1097/GOX.000000000000104

Disclosure: *The authors have no financial interest to declare in relation to the content of this article. The Article Processing Charge was paid for by the authors.*