

Acellular Dermal Matrix in Reconstructive Breast Surgery: Survey of Current Practice among Plastic Surgeons

Ahmed M. S. Ibrahim, MD*
 Pieter G. L. Koolen, MD*
 Azra A. Ashraf, MD, MPH*
 Kuylhee Kim, MD*
 Marc A. M. Mureau, MD,
 PhD†
 Bernard T. Lee, MD, MBA*
 Samuel J. Lin, MD*

Background: Acellular dermal matrices (ADMs) in plastic surgery have become increasingly popular particularly for breast reconstruction. Despite their advantages, questions exist regarding their association with a possible increased incidence of complications. We describe a collective experience of plastic surgeons' use of ADMs in reconstructive breast surgery using an internet-based survey.

Methods: Members of the American Society of Plastic Surgeons were recruited through voluntary, anonymous participation in an online survey. The web-based survey garnered information about participant demographics and their experience with ADM use in breast reconstruction procedures. After responses were collected, all data were anonymously processed.

Results: Data were ascertained through 365 physician responses of which 99% ($n = 361$) completed the survey. The majority of participants were men (84.5%) between 51 and 60 years (37.4%); 84.2% used ADM in breast reconstruction, including radiated patients (79.7%). ADM use was not favored for nipple reconstruction (81.5%); 94.6% of participants used drains, and 87.8% administered antibiotics postoperatively. The most common complications were seroma (70.9%) and infection (16%), although 57.4% claimed anecdotally that overall complication rate was unchanged after incorporating ADM into their practice. High cost was a deterrent for ADM use (37.5%).

Conclusions: Plastic surgeons currently use ADM in breast reconstruction for both immediate and staged procedures. Of those responding, a majority of plastic surgeons will incorporate drains and use postoperative antibiotics for more than 48 hours. (*Plast Reconstr Surg Glob Open* 2015;3:e381; doi: 10.1097/GOX.0000000000000148; Published online 24 April 2015.)

Acellular dermal matrices (ADMs) in plastic surgery have become increasingly popular.¹ The use of ADMs together with improved techniques has helped to solve surgical problems lacking simple surgical solutions.² Not only do these biologic

meshes provide increased structural strength but also they promote rapid vascular ingrowth potentially serving as a scaffold for formation of new tissue.³ Since their availability in the 1990s, the list of indications for their use continues to grow.¹ Among other indications, ADMs have been incorporated into abdominal wall reconstruction, extremity surgery, eyelid reconstruction, and nasal reconstruction.⁴⁻⁷

From the *Division of Plastic Surgery, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Mass.; and †Department of Plastic and Reconstructive Surgery, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands.

Received for publication June 13, 2014; accepted March 30, 2015.

Drs. Ibrahim and Koolen are co-first authors.

Disclaimer: The web-based survey site <http://www.surveymonkey.com> was used to obtain the necessary information. Data extrapolation, statistical analysis, and conclusions reached are the result of the work done by authors of this study.

Copyright © 2015 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of The American Society of Plastic Surgeons. All rights reserved. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 3.0 License, where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially.

DOI: 10.1097/GOX.0000000000000148